



Subinhibitory concentrations of antibiotics affect biofilm formation in *Listeria monocytogenes*

Knudsen, Gitte Maegaard; Gram, Lone

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Knudsen, G. M., & Gram, L. (2012). *Subinhibitory concentrations of antibiotics affect biofilm formation in Listeria monocytogenes*. Poster session presented at ASM 112th General Meeting (ASM 2012), San Francisco, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Subinhibitory concentrations of antibiotics affect biofilm formation in *Listeria monocytogenes*

Gitte M. Knudsen (gmkn@food.dtu.dk) and Lone Gram



Background

The Gram-positive bacteria *Listeria monocytogenes* has a pathogenic and saprophytic life-style and can be exposed to subinhibitory antibiotic concentrations in both niches. These concentrations affect gene expression and physiology such as biofilm formation, cell morphology etc. in several pathogens¹. We have previously found that subinhibitory antibiotic concentrations alter stress and virulence gene expression and cause enhanced stress sensitivity but do not affect Caco-2 cell invasion². The effect of conventional antibiotics on biofilm formation is unknown although several studies have investigated effect of novel antimicrobials.

Purpose

The aim of this study was to determine whether subinhibitory antibiotic concentrations affect *L. monocytogenes* biofilm formation and if so, determine potential mechanisms.

Methods and Materials

Biofilm formation was investigated in BHI broth at 30 and 37°C and quantified using crystal violet (O'Toole & Kolter assay). Bacterial counts were determined of the planktonic part of the culture.

Swimming motility was investigated when exposed to 2-fold concentration of antibiotics in BHI with 0.25% agar at 25 and 37°C.

Biofilm formation

Subinhibitory concentration of ampicillin and gentamicin significantly reduced the biofilm formation at 37°C (Fig. 1A) whereas the same antibiotic increased the biofilm formation at 30°C (Fig. 1B). At both 30 and 37°C, co-trimoxazole enhanced biofilm formation even though the bacterial count was reduced.




Fig.1. Subinhibitory antibiotic concentration change the biofilm formation at 37°C (A) and 30°C (B). Asterisk denote $P < 0.05$ when comparing biofilm of the control to the antibiotic exposed biofilm. Red bar indicate that bacterial count was significantly reduced as compared to control.


Morphology

Subinhibitory concentrations of ampicillin and co-trimoxazole, but not gentamicin and tetracycline, caused minor changes to morphology of the planktonic cells.

Motility

Compared to the control, reduced swimming diameter was observed for all four antibiotics (Fig. 2) however, the bacterial count per plate was also reduced. Thus, the effect was caused by growth reduction of *L. monocytogenes* and not reduced motility.

Fig.2. Swimming at 25°C in BHI with 0.25% agar with 0.125 µg/ml ampicillin, 0.125 µg/ml tetracycline, 8 µg/ml gentamicin and 0.25 µg/ml co-trimoxazole.



Conclusion and Perspective

Subinhibitory concentration of four antibiotics affect *L. monocytogenes* biofilm formation and the antibiotic effect was temperature-dependent. Subinhibitory antibiotic concentrations did not change **motility** and caused only minor modification of the *L. monocytogenes* **morphology** indicating that the altered biofilm capacity was caused by other mechanisms such as c-di-GMP inactivation³, eDNA⁴ or AI-2/LuxS system⁵.

The **temperatur-dependent effect** of the antibiotics on biofilm formation could be advantageous for the organism as enhanced biofilm formation at temperatures below body temperature might **facilitate persistence** in the environment.

Acknowledgement

The work was supported by the Danish Council for Independent Research, Technology and Production Sciences (FTP) grant 09-066098 (274-08-0531).

References

- 1) Romero *et al.* (2011) Chem. Rev. 111:5492
- 2) Knudsen *et al.* (2012) Submitted to JAM
- 3) Hoffman *et al.* (2005) Nature 436: 1171
- 4) Hsu *et al.* (2011) FEMS Immunol. Med. Microbiol 63:236
- 5) Ahmed *et al.* (2009) Antimicrob. Agents Chemother. 53: 4258

The Bacterial Ecophysiology and Biotechnology Group, National Food Institute, Technical University of Denmark, Soeltofts Plads Bldg. 221, DK-2800 Kgs. Lyngby, Denmark